arXiv:1702.08815v1 [math.DS] 28 Feb 2017

INFINITELY MANY PERIODIC ORBITS JUST ABOVE THE MAÑÉ CRITICAL VALUE ON THE 2-SPHERE

GABRIELE BENEDETTI AND MARCO MAZZUCCHELLI

To the memory of John Norman Mather (1942–2017)

ABSTRACT. We introduce a new critical value $c_\infty(L)$ for Tonelli Lagrangians L on the tangent bundle of the 2-sphere without minimizing measures supported on a point. We show that $c_\infty(L)$ is strictly larger than the Mañé critical value $c(L)$, and on every energy level $e \in (c(L), c_\infty(L))$ there exist infinitely many periodic orbits of the Lagrangian system of L, one of which is a local minimizer of the free-period action functional. This has applications to Finsler metrics of Randers type on the 2-sphere. We show that, under a suitable criticality assumption on a given Randers metric, after rescaling its magnetic part with a sufficiently large multiplicative constant, the new metric admits infinitely many closed geodesics, one of which is a waist. Examples of critical Randers metrics include the celebrated Katok metric.

CONTENTS

1. Introduction and main results 1
2. Some remarks on Tonelli Lagrangians 6
3. A minimizer at the Mañé critical value 8
4. Periodic orbits just above the Mañé critical value 9
5. A sufficient condition for the inequality $e_0(L) < c_u(L)$ 12
6. Criticality of the Katok metric 14
References 17

1. Introduction and main results

In this paper, we continue the study started in [AMP15, AMMP17, AAB+17] of the periodic orbits problem for Tonelli Lagrangian systems on the 2-sphere. Such systems are defined by a Tonelli Lagrangian $L : \mathrm{T}S^2 \to \mathbb{R}$, which is a smooth function whose restriction to any fiber of the tangent bundle $\mathrm{T}S^2$ is superlinear with positive definite Hessian. Examples of Tonelli Lagrangians that arise naturally in classical mechanics are

$$L_{g,\theta,V} : \mathrm{T}S^2 \to \mathbb{R}, \qquad L_{g,\theta,V}(q,v) = \tfrac{1}{2} g_q(v,v) + \theta_q(v) - V(q), \tag{1.1}$$

where g is a Riemannian metric, θ a 1-form, and V a function on S^2. When the function V vanishes, we say that the Lagrangian $L_{g,\theta} = L_{g,\theta,0}$ is **magnetic** since it

Date: February 28, 2017.
2000 *Mathematics Subject Classification.* 37J45, 58E05.
Key words and phrases. Tonelli Lagrangians, magnetic flows, periodic orbits, Mañé critical values, Randers metrics, Katok's metrics, closed geodesics.

models the motion of a charged particle under the effect of a stationary magnetic field. As we will observe later in this introduction, magnetic Lagrangians are of importance also in Finsler geometry, since they yield a one parameter family of Randers metrics. We refer the reader to [Fat08, CI99, Abb13, Sor15] for more details on the Tonelli setting in Aubry-Mather theory, weak KAM theory, and symplectic dynamics.

The tangent bundle $\mathrm{T}S^2$ is partitioned into (possibly singular) energy hypersurfaces $E^{-1}(e)$, where E is the energy function

$$E : \mathrm{T}S^2 \to \mathbb{R}, \qquad E(q,v) = \partial_v L(q,v)v - L(q,v).$$

On each energy hypersurface, L defines an Euler-Lagrange flow

$$\phi_L^t : E^{-1}(e) \to E^{-1}(e), \qquad \phi_L^t(\gamma(0),\dot\gamma(0)) = (\gamma(t),\dot\gamma(t)),$$

where the curve $\gamma : \mathbb{R} \to M$ is a smooth solution of the Euler-Lagrange equation

$$\tfrac{\mathrm{d}}{\mathrm{d}t}\partial_v L(\gamma(t),\dot\gamma(t)) - \partial_q L(\gamma(t),\dot\gamma(t)) = 0. \tag{1.2}$$

On a 2-sphere two energy values are known to mark important changes in the qualitative properties of the dynamics. The first one is

$$e_0(L) := \max_{q\in S^2} E(q,0),$$

which is also the maximal $e \in \mathbb{R}$ such that the energy level $E^{-1}(e)$ contains a constant solution of the Lagrangian system (1.2).

The Lagrangian dynamics is variational, meaning that the orbits of the Euler-Lagrange flow can be characterized as critical points of a suitable action functional. Given a periodic curve $\gamma : \mathbb{R}/p\mathbb{Z} \to S^2$, its action at energy e is the quantity

$$\mathcal{S}_e(\gamma) = \int_0^p L(\gamma(t),\dot\gamma(t))\,\mathrm{d}t + pe.$$

The functional $\mathcal{S}_e$, which is defined on the space of periodic curves on M with any possible period, is called the free-period action with energy e. Formally, $\mathcal{S}_e$ is a lower semicontinuous functional of the form

$$\mathcal{S}_e : \Lambda S^2 \times (0,\infty) \to \mathbb{R} \cup \{+\infty\},$$

where $\Lambda S^2 = W^{1,2}(\mathbb{R}/\mathbb{Z}, S^2)$ is a Sobolev manifold of 1-periodic curves, and we identify a pair $(\Gamma, p) \in \Lambda S^2 \times (0,\infty)$ with the p-periodic curve $\gamma(t) = \Gamma(t/p)$. The periodic orbits γ of the Lagrangian system (1.2) with energy $E(\gamma(t),\dot\gamma(t)) \equiv e$ are precisely the critical points of $\mathcal{S}_e$. The second relevant energy level for the dynamics is the Mañé critical value

$$c(L) := \min\big\{e \in \mathbb{R} \;\big|\; \mathcal{S}_e \geq 0\big\}.$$

We refer the reader to, e.g., [Mat91, Mañ97, CI99, Abb13] for a comprehensive introduction to the Mañé critical value.

A recent result due to Asselle and the second author [AM16, Theorem 1.1], first proved by Taimanov [Tai91, Tai92] and Contreras, Macarini, and Paternain [CMP04] in the special case of magnetic Lagrangians, asserts that for every energy value $e \in (e_0(L), c(L))$ there exists a periodic orbit that is a local minimizer of $\mathcal{S}_e$. Thus, it is natural to ask whether local minimizers exist also for energies $e \geq c(L)$. If $e_0(L) = c(L)$ there exists a global minimizer that is a constant curve at level $c(L)$. Indeed, if $\gamma_q : \mathbb{R}/p\mathbb{Z} \to M$ is the curve $\gamma_q(t) \equiv q$, its action is given by

$$\mathcal{S}_e(\gamma_q) = p\,(L(q,0) + e) = p\,(e - E(q,0)) \tag{1.3}$$

and, if $E(q,0) = e_0(L)$, we conclude that $\mathcal{S}_{e_0(L)}(\gamma_q) = 0$. Hence, if $e_0(L) = c(L)$, the functional $\mathcal{S}_{e_0(L)}$ is non-negative, and γ_q is a global minimizer. The next simple example shows that, when $e_0(L) = c(L)$, all the minimizers of $\mathcal{S}_{c(L)}$ may actually be constant, and $\mathcal{S}_e$ may not admit any minimizer for $e > c(L)$.

Example 1.1. Consider the round Riemannian metric g on S^2, and the Lagrangian $L : \mathrm{T}S^2 \to \mathbb{R}$ given by $L(q,v) = \frac{1}{2} g_q(v,v)$. The associated energy function is $E(q,v) = L(q,v)$, and we have $e_0(L) = c(L) = 0$. On $E^{-1}(0)$ all orbits are constant, whereas on any $E^{-1}(e)$ with $e > 0$ the Euler-Lagrange flow is the geodesic flow of the round metric. In particular, each orbit of this flow is periodic and is not a local minimizer of $\mathcal{S}_e$. ■

As a consequence of a result of Haefliger on minimal measures with rational rotation number [CMP04, Prop. 2.1], magnetic Lagrangians with $e_0(L) < c(L)$ possess a periodic orbit at energy $c(L)$ that is a global minimizer of $\mathcal{S}_{c(L)}$. We generalize hereby this result to general Tonelli Lagrangians. As usual, we say that a periodic curve $\gamma : \mathbb{R}/p\mathbb{Z} \to M$ is simple when it has no self-intersections, meaning that $\gamma(t) \neq \gamma(s)$ for all distinct $t, s \in \mathbb{R}/p\mathbb{Z}$.

Theorem 1.2. *Let $L : \mathrm{T}S^2 \to \mathbb{R}$ be a Tonelli Lagrangian such that $e_0(L) < c(L)$. The Lagrangian system of L admits a simple periodic orbit γ with energy $c(L)$ that is a global minimizer of the free-period action functional $\mathcal{S}_{c(L)}$ and satisfies $\mathcal{S}_{c(L)}(\gamma) = 0$.*

The global minimizer γ of $\mathcal{S}_{c(L)}$ is not necessarily unique. Nevertheless, the assumption $e_0(L) < c(L)$ implies that no orbit with energy $c(L)$ is constant, and therefore that the global minimizers of $\mathcal{S}_{c(L)}$ form a compact set. As we will show, this forces the existence of local minimizers of $\mathcal{S}_e$ for all energies e slightly above the Mañé critical value.

Theorem 1.3. *Let $L : \mathrm{T}S^2 \to \mathbb{R}$ be a Tonelli Lagrangian such that $e_0(L) < c(L)$. There exists $\epsilon > 0$ such that, for all energy levels $e \in (c(L), c(L)+\epsilon)$, the Lagrangian system of L admits a simple periodic orbit with energy e that is a local minimizer of the free-period action functional $\mathcal{S}_e$.*

We wish to stress that the minimizers of $\mathcal{S}_e$, for $e > c(L)$, are never global ones. Indeed, Equation (1.3) implies that

$$\inf \mathcal{S}_e = 0, \qquad \forall e \geq c(L). \tag{1.4}$$

If $\gamma = (\Gamma, p)$ were a global minimizer of $\mathcal{S}_e$ with $e > c(L)$, then $\mathcal{S}_e(\gamma) = 0$ and

$$\mathcal{S}_{c(L)}(\gamma) = \mathcal{S}_e(\gamma) + (c(L) - e)p = (c(L) - e)p < 0,$$

contradicting the definition of the Mañé critical value. Theorem 1.3 motivates the definition of a new critical energy value.

Definition 1.4. We define the **critical value** $c_\infty(L)$ to be the supremum of the energy levels $e' > e_0(L)$ such that, for all $e \in (e_0(L), e']$, the free-period action functional $\mathcal{S}_e$ admits a local minimizer.

Remark 1.5. In general, $c_\infty(L)$ is finite, as can be seen in the example of the magnetic Lagrangians associated to the Katok metrics, see Section 6. ■

If $e_0(L) < c(L)$, combining the above mentioned [AM16, Theorem 1.1] with Theorems 1.2 and 1.3 we get the strict inequality

$$c_\infty(L) > c(L). \tag{1.5}$$

The waist theorem for Tonelli Lagrangians on surfaces, which was stated without proof in [AMMP17, Corollary 2.7], asserts that the presence of a local minimizer γ of the free-period action functional $\mathcal{S}_e$ implies the existence of infinitely many other periodic orbits with energy e provided $\{\gamma\}$ is not the whole set of global minimizers of $\mathcal{S}_e$, $\mathcal{S}_e(\gamma) \neq 0$, and $\mathcal{S}_e$ satisfies the Palais-Smale condition. Therefore, as a consequence of this waist theorem and of inequality (1.5), we obtain a new multiplicity result for Tonelli periodic orbits on S^2 at energy levels above the Mañé critical value.

Theorem 1.6. *Let $L : \mathrm{T}S^2 \to \mathbb{R}$ be a Tonelli Lagrangian such that $e_0(L) < c(L)$. For all energy levels $e \in (c(L), c_\infty(L))$, the Lagrangian system of L admits infinitely many periodic orbits with energy e.*

When $L = L_{g,\theta}$ is a magnetic Lagrangian, the condition $e_0(L) < c(L)$ is equivalent to the fact that the 1-form θ is not exact. For more general Tonelli Lagrangians $L : \mathrm{T}S^2 \to \mathbb{R}$, for example if L is given by (1.1) with $V \neq 0$, we provide a sufficient condition for the inequality to hold. The statement is a special case of Proposition 5.1, which is valid for Tonelli Lagrangians on any configuration space.

Proposition 1.7. *Let $L : \mathrm{T}S^2 \to \mathbb{R}$ be a Tonelli Lagrangian. We set $V(q) := L(q, 0)$, $\theta_q := \partial_v L(q, 0)$, $g_q := \partial^2_{vv} L(q, 0)$, and*

$$\lambda : V^{-1}(e_0(L)) \to \mathbb{R}, \qquad \lambda(q) := 2|\mathrm{d}^2 V(q)|^{1/2} - |\mathrm{d}\theta_q|,$$

where $\mathrm{d}^2 V$ denotes the Hessian of V, and the norms are taken with respect to the Riemannian metric g. If the function λ is somewhere negative, then $e_0(L) < c(L)$.

It is well known that for every $e > c(L)$, the Euler-Lagrange flow of L on the energy hypersurface $E^{-1}(e)$ is orbitally equivalent to the geodesic flow of a Finsler metric on the unit tangent bundle of S^2, see [CIPP98, Cor. 2]. For magnetic Lagrangians $L = L_{g,\theta}$, this equivalence is particularly explicit, as we now recall following [Pat99]. Let $\sim$ be the equivalence relation among 1-forms on S^2 such that $\theta' \sim \theta$ if and only if $\theta - \theta'$ is exact. We will write $[\theta]$ for the equivalence class of θ. The Euler-Lagrange flow and the free-period action functional $\mathcal{S}_e$ depend only on the equivalence class $[\theta]$. In particular, the same is true for the critical values $c(L)$ and $c_\infty(L)$. We denote by $|\cdot|$ the norm of tangent and cotangent vectors associated to the Riemannian metric g on S^2. If θ' is a 1-form, we will write its L^∞-norm by

$$\|\theta'\|_\infty := \max\big\{|\theta'_q| \;\big|\; q \in S^2\big\}.$$

The energy function associated to L is given by $E(q, v) = \frac{1}{2}|v|^2$, and thus $e_0(L) = 0$. The Tonelli Hamiltonian $H : \mathrm{T}^*S^2 \to \mathbb{R}$ dual to L is given by

$$H(q, p) = \tfrac{1}{2}|p - \theta_q|^2.$$

By [CIPP98, Theorem A], the Mañé critical value can be computed as a minimax of the Hamiltonian over exact Lagrangian graphs. For the magnetic Lagrangian L, this recipe gives

$$c(L) = \inf_{u \in C^\infty(S^2)} \max_{q \in S^2} H(q, \mathrm{d}u(q)) = \inf_{\theta' \sim \theta} \tfrac{1}{2}\|\theta'\|^2_\infty.$$

Proposition 1.7 implies that $c(L) > e_0(L)$ provided θ is not exact. For all $r > \|\theta\|_\infty$, we denote by F_r the Finsler metric of Randers type

$$F_r(q,v) = r|v| + \theta_q(v).$$

The diffeomorphism

$$\psi_r : E^{-1}(r^2/2) \to F_r^{-1}(1), \qquad \psi_r(q,v) = (q, (r^2 + \theta_q(v))^{-1} v)$$

realizes an orbit equivalence between the Euler-Lagrange flow of L and the geodesic flow of F_r on the respective energy hypersurfaces [Pat99, Lemma 2.1]. We define

$$r(g,[\theta]) := \sqrt{2\,c(L)} = \inf_{\theta' \sim \theta} \|\theta'\|_\infty, \tag{1.6}$$
$$r_\infty(g,[\theta]) := \sqrt{2\,c_\infty(L)}.$$

A periodic curve $\Gamma : \mathbb{R}/\mathbb{Z} \to S^2$ is called a **waist** of the Finsler metric F_r when it is a local minimizer of the length function

$$\mathcal{L}_r : C^\infty(\mathbb{R}/\mathbb{Z}, S^2) \to \mathbb{R}, \qquad \mathcal{L}_r(\Gamma) = \int_0^1 F_r(\Gamma(t), \dot\Gamma(t))\,\mathrm{d}t.$$

The reparametrization of a waist with constant F_r-speed is a closed geodesic for the Finsler metric F_r. Notice that, if $\gamma = (\Gamma, p)$ is a smooth periodic curve with constant energy $E(\gamma(t), \dot\gamma(t)) \equiv \frac{1}{2} r^2$, then $\mathcal{S}_{r^2/2}(\gamma) = \mathcal{L}_r(\Gamma)$. This, together with the fact that $\mathcal{L}_r(\Gamma)$ is independent of the parametrization of Γ, implies that Γ is a waist of F_r whenever $\gamma = (\Gamma, p)$ is a local minimizer of the free-period action functional $\mathcal{S}_{r^2/2}$. Therefore, for magnetic Lagrangians $L = L_{g,\theta}$, Theorems 1.3 and 1.6 can be rephrased as follows.

Theorem 1.8. *Let g be a Riemannian metric and θ a non-exact 1-form on S^2 such that $\|\theta\|_\infty < r_\infty(g,[\theta])$. For all $r \in (\|\theta\|_\infty, r_\infty(g,[\theta]))$ the Finsler metric*

$$F_r(q,v) = r g_q(v,v)^{1/2} + \theta_q(v)$$

has a waist and infinitely many closed geodesics. ■

Given an equivalence class for the relation $\sim$, we can always find a representative θ such that $\|\theta\|_\infty < r_\infty(g,[\theta])$. Therefore, given the geodesic flow of a Randers metric on S^2, we can always find a pair (g,θ) that defines a Randers metric with the same geodesic flow and at the same time satisfies the assumption of Theorem 1.8. Sometimes, there even exist pairs (g,θ) that are in some sense optimal.

Definition 1.9. We say that (g,θ), where g is a Riemannian metric on S^2 and θ a 1-form on S^2, is a **critical pair** if $r(g,[\theta]) = \|\theta\|_\infty$. When $r(g,[\theta]) = \|\theta\|_\infty < 1$, we will also say that the associated metric $F(q,v) = g_q(v,v)^{1/2} + \theta_q(v)$ is critical. ■

Since $r(g,[\theta]) < r_\infty(g,[\theta])$, Theorem 1.8 is always applicable to critical pairs. In Section 6, we will provide a sufficient condition for a pair (g,θ) to be critical. This condition holds, for instance, if (g,θ) is rotationally symmetric (see Remark 6.3). In particular, the celebrated Katok metric on S^2 is critical.

1.1. **Organization of the paper.** In Section 2 we provide two technical statements that are needed in Sections 3 and 4. In Section 3 we prove Theorem 1.2. In Section 4, we prove Theorems 1.3 and 1.6. In Section 5 we prove a more general version of Proposition 1.7. Finally, in Section 6, we show that the Katok metric on S^2 is critical, and we estimate its quantity r_∞.

1.2. **Acknowledgments.** G.B. thanks Luca Asselle for several inspiring conversations about the topics of this paper. M.M. is grateful to Albert Fathi for a fruitful discussion about the proof of Theorem 1.2. M.M. is partially supported by the ANR COSPIN (ANR-13-JS01-0008-01).

2. Some remarks on Tonelli Lagrangians

Let M be a closed manifold, and $L : \mathrm{T}M \to \mathbb{R}$ a Tonelli Lagrangian with energy function $E : \mathrm{T}M \to \mathbb{R}$. Let $\phi_L^t : \mathrm{T}M \to \mathrm{T}M$ denote the Euler-Lagrange flow of L on the whole tangent bundle, and $\pi : \mathrm{T}M \to M$ the base projection.

2.1. **On simple periodic orbits.** By [Fat08, Theorem 2.7.4], for each energy value $e > e_0(L)$ there exists $\tau = \tau(e) > 0$ small enough such that, for all $q \in M$ and $t \in (0, \tau)$, the map $\pi \circ \phi_L^t|_{\mathrm{T}_q M \cap E^{-1}(-\infty, e)}$ is a diffeomorphism onto an open neighborhood of q. Moreover, by [Fat08, Theorem 3.6.1], for all $(q, v) \in M$ with $E(q, v) < e$ and $\sigma \in (0, \tau)$, the curve

$$\gamma : [0, \sigma] \to M, \qquad \gamma(t) = \pi \circ \phi_L^t(q, v)$$

is a strict action minimizer: if $\zeta : [0, \sigma] \to M$ is any other absolutely continuous curve such that $\zeta(0) = \gamma(0)$ and $\zeta(\sigma) = \gamma(\sigma)$, then

$$\int_0^\sigma L(\gamma(t), \dot\gamma(t))\, \mathrm{d}t < \int_0^\sigma L(\zeta(t), \dot\zeta(t))\, \mathrm{d}t. \tag{2.1}$$

The following remark will be needed in order to show that the the periodic orbits provided by Theorem 1.3 are simple.

Lemma 2.1. *Let M be a closed manifold, and $L : \mathrm{T}M \to \mathbb{R}$ a Tonelli Lagrangian. For every simple periodic orbit $\gamma = (\Gamma, p) \in W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times (0, \infty)$ of the Lagrangian system of L, there exists a neighborhood $\mathcal{U} \subset W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times (0, \infty)$ such that any periodic orbit of the Lagrangian system of L in $\mathcal{U}$ is simple as well.*

Proof. Let $\gamma_n = (\Gamma_n, p_n)$ be a sequence of periodic orbits that converges to the simple one $\gamma = (\Gamma, p)$ in $W^{1,2}(\mathbb{R}/\mathbb{Z}, M) \times (0, \infty)$ as $n \to \infty$. Let $e_n := E(\gamma_n(0), \dot\gamma_n(0))$. We claim that there exists $e \in \mathbb{R}$ such that $e_n < e$ for all $n \in \mathbb{N}$. Otherwise, up to passing to a subsequence, we would have $e_n \to \infty$. This would imply that $|\dot\gamma_n(t)| \to \infty$ for all $t \in \mathbb{R}$, where $|\cdot|$ denotes an arbitrary Riemannian norm. Therefore, since the periods p_n are bounded away from zero, we would obtain that

$$\mathrm{length}(\gamma_n) = \int_0^{p_n} |\dot\gamma_n(t)|\, \mathrm{d}t \to \infty,$$

which is impossible, for $\mathrm{length}(\gamma_n) \to \mathrm{length}(\gamma)$.

Let $\tau = \tau(e) > 0$ be the constant introduced above, so that for all $q \in M$ and $t \in (0, \tau)$ the map $\pi \circ \phi_L^t|_{\mathrm{T}_q M \cap E^{-1}(-\infty, e)}$ is a diffeomorphism onto an open neighborhood of q. Since, for all $t \in (0, \tau)$, we have that $\gamma_n(t) = \pi \circ \phi_L^t(\gamma_n(0), \dot\gamma_n(0)) \to \gamma(t)$ as $n \to \infty$, we infer that $\dot\gamma_n(0) \to \dot\gamma(0)$. Therefore, Γ_n converges to Γ in $C^\infty(\mathbb{R}/\mathbb{Z}, M)$. Since being a simple curve is a C^1-open condition, we conclude that, for n large enough, γ_n is a simple curve. ∎

2.2. **Action minimizers and modifications of the Lagrangian.** Let

$$\mathcal{S}_e : \Lambda M \times (0,\infty) \to \mathbb{R} \cup \{\infty\},$$
$$\mathcal{S}_e(\gamma) = \int_0^p L(\gamma(t), \dot\gamma(t))\,\mathrm{d}t + pe, \qquad \forall \gamma = (\Gamma, p) \in \Lambda M \times (0,\infty).$$

be the free-period action functional associated to L at energy e. Any local minimizer γ of $\mathcal{S}_e$ is a periodic orbits with energy e. If we modify the Lagrangian away from $E^{-1}(e)$, γ remains a periodic orbit of the new Lagrangian system, but a priori it may not be a minimizer of the new free-period action functional. However, we have the following statement.

Lemma 2.2. *Let $L' : \mathrm{T}M \to \mathbb{R}$ be a Tonelli Lagrangian that coincide with L on the sublevel set $E^{-1}(-\infty, e']$, for some $e' > e$, and let $\mathcal{S}'_e$ be the free-period action functional of L'. A local minimizer of $\mathcal{S}'_e$ is a local minimizer of $\mathcal{S}_e$ as well.*

Remark 2.3. The lemma becomes trivial if one considers the free-period action functionals defined on the space of C^1 periodic curves. However, in order to apply global methods from nonlinear analysis, it is more suitable to work with the Sobolev loop space $\Lambda M = W^{1/2}(\mathbb{R}/\mathbb{Z}, M)$. ■

Proof of Lemma 2.2. Assume by contradiction that there exists a local minimizer $\gamma = (\Gamma, p)$ of $\mathcal{S}'_e$ that is not a local minimizer of $\mathcal{S}_e$. Therefore, there exists a sequence $\zeta_n = (Z_n, p_n) \in \Lambda M \times (0,\infty)$ such that $\mathcal{S}_e(\zeta_n) < \mathcal{S}_e(\gamma)$ and $\zeta_n \to \gamma$ in $\Lambda M \times (0,\infty)$. Consider the quantity $\tau = \tau(e) > 0$ introduced above, and $k \in \mathbb{N}$ large enough so that $p/k < \tau$. For all n large enough, we have that $p_n/k < \tau$ and, for each $t \in \mathbb{R}/p_n\mathbb{Z}$, the map $\psi_{n,t} : \mathrm{T}_{\zeta_n(t)}M \cap E^{-1}(-\infty, e) \to M$ given by

$$\psi_{n,t}(v) = \pi \circ \phi_L^{p_n/k}(\zeta_n(t), v)$$

is a diffeomorphism onto a neighborhood of $\{\zeta_n(t), \zeta_n(t + p_n/k)\}$. We set

$$v_{n,i} := \psi^{-1}_{n,p_n i/k}\big(\zeta_n\big(\tfrac{(i+1)p_n}{k}\big)\big), \qquad i = 0, ..., k-1,$$

and we define the periodic curve $\gamma_n = (\Gamma_n, p_n) \in \Lambda M \times (0,\infty)$ by

$$\gamma_n(\tfrac{ip_n}{k} + t) := \pi \circ \phi_L^t\big(\zeta_n\big(\tfrac{ip_n}{k}\big), v_{n,i}\big).$$

Notice that γ_n is a continuous and piecewise broken periodic orbit of the Lagrangian system of L satisfying $\mathcal{S}_k(\gamma_n) < \mathcal{S}_k(\zeta_n)$. Moreover, since $\gamma_n(\frac{ip_n}{k}) = \zeta_n(\frac{ip_n}{k}) \to \gamma(\frac{ip}{k})$ as $n \to \infty$, we have that $\Gamma_n|_{[i/k,(i+1)/k]} \to \Gamma|_{[i/k,(i+1)/k]}$ in the C^∞ topology. In particular, for all $n \in \mathbb{N}$ large enough and $t \in \mathbb{R}/p_n\mathbb{Z}$, we have that $E(\gamma_n(t), \dot\gamma_n(t^+)) < e'$, which implies

$$\mathcal{S}_e(\gamma_n) = \mathcal{S}'_e(\gamma_n) < \mathcal{S}'_e(\gamma) = \mathcal{S}_e(\gamma).$$

This contradicts the fact that γ is a minimizer of $\mathcal{S}_e$. ■

2.3. **The Tonelli free-period action functional.** The free-period action functional $\mathcal{S}_e$ of a Tonelli Lagrangian L does not have good functional properties when L is not quadratic at infinity: it is not continuous, and its sublevel sets do not enjoy any sort of weak compactness. However, we can circumvent these issues by modifying the Lagrangian L away from the region where we consider its dynamics. Since this paper deals with periodic orbits with energy equal to or just above the Mañé critical value $c(L)$, we can focus on the energy range $[\min E, c(L)+1]$. We can thus replace L by a Tonelli Lagrangian $L' : \mathrm{T}M \to \mathbb{R}$ that coincides with L in the energy sublevel

set $E^{-1}(-\infty, c(L)+1]$, and is quadratic at infinity. This latter condition means that there exists a compact set $Q \subset \mathrm{T}M$ such that, for any $q \in M$, the function $L'(q,\cdot)$ coincides with a polynomial of degree 2 outside $Q \cap \mathrm{T}_q M$. A construction of the Lagrangian L' can be found in, e.g., [CIPP00, Prop. 18]. By [CIPP00, Lemma 19], we have $c(L) = c(L')$. Moreover, the free-period action functional $\mathcal{S}'_e$ associated to L' is well behaved: it is $C^{1,1}$, and satisfies the Palais-Smale condition on subsets of the form $\Lambda M \times [p_-, p_+]$, for any compact interval $[p_-, p_+] \subset (0,\infty)$, see [Abb13, Lemmas 3.1 and 5.3]. By Lemma 2.2, for all $e \in [\min E, c(L)+1]$, the local minimizers of $\mathcal{S}_e$ and $\mathcal{S}'_e$ coincide.

Thanks to this remark, throughout Sections 3 and 4, while working in the energy range $[\min E, c(L)+1]$, we will be able to assume without loss of generality that the Tonelli Lagrangian $L : \mathrm{T}S^2 \to \mathbb{R}$ considered there is quadratic at infinity, so that the associated free-period action is a well behaved functional of the form $\mathcal{S}_e : \Lambda M \times (0,\infty) \to \mathbb{R}$, for $e \in [\min E, c(L)+1]$.

3. A minimizer at the Mañé critical value

Let $L : \mathrm{T}S^2 \to \mathbb{R}$ be a Tonelli Lagrangian with associated free-period action functionals $\mathcal{S}_e$, $e \in \mathbb{R}$. We say that a periodic curve $\gamma = (\Gamma, p) \in \Lambda S^2 \times (0,\infty)$ is non-iterated when p is its minimal period (that is, 1 is the minimal period of Γ).

Lemma 3.1. *Any non-iterated global minimizer $\gamma = (\Gamma, p)$ of the free-period action functional $\mathcal{S}_{c(L)}$ is a simple periodic orbit with action $\mathcal{S}_{c(L)}(\gamma) = 0$.*

Proof. Let $\gamma = (\Gamma, p)$ be a non-iterated global minimizer of $\mathcal{S}_{c(L)}$. Being a critical point of $\mathcal{S}_{c(L)}$, γ is a periodic orbit with energy $c(L)$. By (1.4), we have $\mathcal{S}_{c(L)}(\gamma) = 0$. Assume by contradiction that there exist $t_0, t_1 \in [0,p]$ such that $p_0 := t_1 - t_0 \in (0,p)$ and $\gamma(t_0) = \gamma(t_1)$. We define the loops $\gamma_0 : \mathbb{R}/p_0\mathbb{Z} \to S^2$, $\gamma_0(t) = \gamma(t_0+t)$ and $\gamma_1 : \mathbb{R}/(p-p_0)\mathbb{Z} \to S^2$, $\gamma_1(t) = \gamma(t_1+t)$. Since $\mathcal{S}_{c(L)}(\gamma) = \mathcal{S}_{c(L)}(\gamma_0) + \mathcal{S}_{c(L)}(\gamma_1)$ and γ is a global minimizer of $\mathcal{S}_{c(L)}$, we must have $\mathcal{S}_{c(L)}(\gamma_0) = \mathcal{S}_{c(L)}(\gamma_1) = 0$. Notice that γ_0 is not smooth at $t = 0$, for otherwise γ would have been an iterated periodic orbit. Therefore, for $\epsilon > 0$ small enough, there exists a smooth curve $\zeta : [0,\delta] \to S^2$ such that $\zeta(0) = \gamma_0(-\epsilon)$, $\zeta(\delta) = \gamma_0(\epsilon)$, and

$$\int_0^\delta L(\zeta(t), \dot\zeta(t))\,\mathrm{d}t + \delta\, c(L) > \int_{-\epsilon}^{\epsilon} L(\gamma_0(t), \dot\gamma_0(t))\,\mathrm{d}t + 2\epsilon\, c(L),$$

see e.g. [AM16, Lemma 2.3(ii)]. We replace the portion $\gamma_0|_{[-\epsilon,\epsilon]}$ of γ_0 with the curve ζ. This produces a piecewise smooth periodic curve $\gamma_2 : \mathbb{R}/(p_0 - 2\epsilon + \delta)\mathbb{Z}$ such that $\mathcal{S}_{c(L)}(\gamma_2) < \mathcal{S}_{c(L)}(\gamma_0) = 0$, which contradicts (1.4). ∎

Before proceeding with the proof of Theorem 1.2 we provide some preliminaries that will be useful also in Section 4. We fix an arbitrary Riemannian metric g on S^2 that we will employ for computing distances and lengths of curves. For each $e' > e_0(L)$, there is a constant $\ell_{\min}(e') > 0$ such that, for all $e \geq e'$, every absolutely continuous periodic curve $\gamma : \mathbb{R}/p\mathbb{Z} \to S^2$ that is entirely contained in a Riemannian ball of diameter $\ell_{\min}(e')$ satisfies $\mathcal{S}_e(\gamma) > 0$, see [AM16, Corollary 2.5]. We denote by θ the 1-form on S^2 given by $\theta_q(v) = \partial_v L(q,0)v$. If $\gamma = (\Gamma, p)$ is a simple curve, a simple application of Stokes' Theorem [AM16, Lemma 2.10] yields the estimate

$$p \leq \frac{\mathcal{S}_e(\gamma) + \int_{S^2} |\mathrm{d}\theta|}{e - e_0(L)}. \tag{3.1}$$

Proof of Theorem 1.2. By [AM16, Theorem 1.1], for every $e \in (e_0(L), c(L))$ there exists a periodic orbit γ_e of the Lagrangian system of L with energy e such that

(i) $\gamma_e = (\Gamma_e, p_e)$ is a local minimizer of $\mathcal{S}_e : \Lambda S^2 \times (0,\infty) \to \mathbb{R}$,
(ii) $\mathcal{S}_e(\gamma_e) < 0$.
(iii) the loop $\gamma_e : \mathbb{R}/p_e\mathbb{Z} \hookrightarrow S^2$ is simple.

Fix now $e' \in (e_0(L), c(L))$. By (ii) and (3.1), for all $e \in [e', c(L))$ we have the upper bounds

$$p_e \le p_{\max} := (e' - e_0(L))^{-1} \int_{S^2} |\mathrm{d}\theta|,$$
$$\mathrm{length}(\gamma_e) \le \ell_{\max} := p_{\max} \max\left\{ g_q(v,v)^{1/2} \;\middle|\; (q,v) \in E^{-1}(c(L)) \right\},$$

and the lower bounds

$$\mathrm{length}(\gamma_e) \ge \ell_{\min} > 0,$$
$$p_e \ge p_{\min} := \ell_{\min} \min\left\{ g_q(v,v)^{-1/2} \;\middle|\; (q,v) \in E^{-1}(c(L)) \right\} > 0,$$

where $\ell_{\min} = \ell_{\min}(e')$. Therefore, we can find a sequence of energy values $e_n \in [e', c(L))$ such that $e_n \to c(L)$ and $\gamma_{e_n} = (\Gamma_{e_n}, p_{e_n}) \to \gamma = (\Gamma, p)$ in $\Lambda S^2 \times (0,\infty)$. The function $(e,\zeta) \mapsto \mathcal{S}_e(\zeta)$ is continuous, and hence $\mathcal{S}_{e_n}(\gamma_{e_n}) \to \mathcal{S}_{c(L)}(\gamma)$. This, together with property (ii), implies that $\mathcal{S}_{c(L)}(\gamma) \le 0$. By (1.4), γ is a global minimizer of $\mathcal{S}_{c(L)}$ with $\mathcal{S}_{c(L)}(\gamma) = 0$. Since γ is the limit of non-self-intersecting periodic curves γ_{e_n} with period $p_{e_n} \to p$, γ must be non-iterated, that is, p must be its minimal period. Lemma 3.1 implies that γ is non-self-intersecting. ∎

4. Periodic orbits just above the Mañé critical value

Given a periodic curve $\gamma = (\Gamma, p) \in \Lambda S^2 \times (0,\infty)$ and a positive integer m, we write $\gamma^m = (\Gamma^m, mp) \in \Lambda S^2 \times (0,\infty)$ for the m-th iterate of γ, where $\Gamma^m(t) = \Gamma(mt)$. We denote by

$$\psi^m : \Lambda S^2 \times (0,\infty) \hookrightarrow \Lambda S^2 \times (0,\infty)$$

the iteration map $\psi^m(\gamma) = \gamma^m$. The free-period action functional $\mathcal{S}_e$ is equivariant with respect to the iteration action, in the sense that $\mathcal{S}_e \circ \psi^m = m\,\mathcal{S}_e$.

Let us assume $e_0(L) < c(L)$ and consider the set of global minimizers

$$\mathcal{K} := \mathcal{S}_{c(L)}^{-1}(0) \subset \Lambda S^2 \times (0,\infty).$$

By Theorem 1.2, $\mathcal{K}$ is non-empty and we denote by $\mathcal{K}_1 \subset \mathcal{K}$ the subset of non-iterated periodic curves in $\mathcal{K}$. Since a periodic curve has zero action if and only if the same is true for any of its iterates, we have $\mathcal{K} := \mathcal{K}_1 \cup \mathcal{K}_2 \cup \mathcal{K}_3 \cup ...$, where $\mathcal{K}_m := \psi^m(\mathcal{K}_1)$. Lemma 3.1 further implies that every periodic curve in $\mathcal{K}_1$ is simple. By (3.1), the periods of the curves $\gamma = (\Gamma, p) \in \mathcal{K}_1$ are uniformly bounded from above as

$$p \le p_{\max} := (c(L) - e_0(L))^{-1} \int_{S^2} |\mathrm{d}\theta|,$$

The periods of curves $\gamma = (\Gamma, p) \in \mathcal{K}_1$ are also uniformly bounded from below as

$$p \ge p_{\min} := \ell_{\min} \max\left\{ |v|_g \;\middle|\; (q,v) \in E^{-1}(c(L)) \right\}^{-1} > 0,$$

where $\ell_{\min} = \ell_{\min}(c(L))$.

Lemma 4.1. *The space $\mathcal{K}_1$ is compact, non-empty, and it admits an open neighborhood $\mathcal{U} \subset \Lambda S^2 \times \left[p_{\min}/2, 2p_{\max}\right]$ such that $\overline{\mathcal{U}} \cap \mathcal{K} = \mathcal{K}_1$.*

Proof. Since the free-period action functional $\mathcal{S}_{c(L)}$ is continuous, $\mathcal{K}_1$ is closed. Let $\gamma_i = (\Gamma_i, p_i) \in \mathcal{K}_1$ be an arbitrary sequence of periodic curves. We already know that $p_{\min} \leq p_i \leq p_{\max}$. The corresponding sequence of the initial velocity vectors $(\gamma_i(0), \dot{\gamma}_i(0))$ is contained into the compact energy hypersurface $E^{-1}(c(L))$. Therefore, up to extracting a subsequence, we see that $\gamma_i \to \gamma = (\Gamma, p) \in \mathrm{crit}(S_{c(L)})$. Since $\mathcal{S}_{c(L)}(\gamma_i) = 0$ for all $i \in \mathbb{N}$, we have that $\mathcal{S}_{c(L)}(\gamma) = 0$, and thus $\gamma \in \mathcal{K}$. As the curves γ_i are non-iterated, the same is true for the limit curve γ. Therefore, γ belongs to $\mathcal{K}_1$. This shows that $\mathcal{K}_1$ is compact.

Since the curves in $\mathcal{K}_1$ are simple, $\mathcal{K}_m$ and $\mathcal{K}_n$ are disjoint if $m \neq n$. In particular, we can find an open neighborhood $\mathcal{U} \subset \Lambda S^2 \times [p_{\min}/2, 2p_{\max}]$ of $\mathcal{K}_1$ whose closure is disjoint from $\mathcal{K}_m$ for all $m \in \{1, ..., \lceil 2p_{\max}/p_{\min} \rceil\}$. Such neighborhood is also disjoint from $\mathcal{K}_m$ for all $m > \lceil 2p_{\max}/p_{\min} \rceil$; indeed, any curve γ contained in one such $\mathcal{K}_m$ is of the form $\gamma = (\Gamma, p)$ with $p \geq mp_{\min} > 2p_{\max}$. ∎

Lemma 4.2. *There exist a number $\epsilon \in (0,1)$ and an arbitrarily small open neighborhood $\mathcal{W} \subseteq \mathcal{U}$ of $\mathcal{K}_1$ such that $\inf \mathcal{S}_e|_{\partial\mathcal{W}} > \inf \mathcal{S}_e|_{\mathcal{W}}$ for all $e \in [c(L), c(L) + \epsilon)$.*

Proof. Let d be the distance on $\Lambda S^2 \times (0, \infty)$ induced by its standard Riemannian metric. For each $r > 0$, we denote by $\mathcal{W}_r$ the open neighborhood of $\mathcal{K}_1$ of radius r, that is

$$\mathcal{W}_r = \big\{\gamma \in \Lambda S^2 \times (0,\infty) \;\big|\; d(\gamma, \zeta) < r \text{ for some } \zeta \in \mathcal{K}_1\big\}.$$

We choose $r > 0$ small enough so that $\mathcal{W}_{3r} \subseteq \mathcal{U}$, and the gradient norm $\|\nabla \mathcal{S}_{c(L)}\|$ is bounded from above on $\mathcal{W}_{3r}$. We set $\mathcal{C}$ to be the set of critical points of $\mathcal{S}_{c(L)}$ contained in the closure of $\mathcal{W}_{3r} \setminus \mathcal{W}_r$. Since $\mathcal{S}_e$ satisfies the Palais-Smale condition on $\Lambda S^2 \times [p_{\min}/2, 2p_{\max}]$, Lemma 4.1 implies that

$$a := \inf \mathcal{S}_{c(L)}|_{\mathcal{C}} > 0.$$

Thus, the closure of $\{\mathcal{S}_{c(L)} < a/2\} \cap (\mathcal{W}_{3r} \setminus \mathcal{W}_r)$ does not contain any critical point of $\mathcal{S}_{c(L)}$ and, again by the Palais-Smale condition, there exists $\delta > 0$ such that $\|\nabla \mathcal{S}_{c(L)}\| \geq \delta$ on $\{\mathcal{S}_{c(L)} < a/2\} \cap (\mathcal{W}_{3r} \setminus \mathcal{W}_r)$.

We set $\mathcal{W} := \mathcal{W}_{2r}$, and we claim that $\inf \mathcal{S}_{c(L)}|_{\partial\mathcal{W}} > \inf \mathcal{S}_{c(L)}|_{\mathcal{W}} = 0$. Indeed, assume by contradiction that there exists a sequence $\{\gamma_i \mid i \in \mathbb{N}\} \subset \partial\mathcal{W}$ with $\mathcal{S}_{c(L)}(\gamma_i) < 1/i$. Fix $0 < b < \min\{a/2, \delta r\}$ and $i > 1/b$. Notice that the curve γ_i is contained in the sublevel set $\{\mathcal{S}_{c(L)} < a/2\}$. Consider the anti-gradient flow Φ_s of the functional $\mathcal{S}_{c(L)}$, that is, the partial flow defined by the ordinary differential equation

$$\tfrac{\mathrm{d}}{\mathrm{d}s}\Phi_s = -\nabla \mathcal{S}_{c(L)} \circ \Phi_s.$$

Since $\|\nabla \mathcal{S}_{c(L)}\|$ is bounded from above on $\mathcal{W}_{3r}$, every orbit $s \mapsto \Phi_s(\gamma)$ that does not stay inside $\mathcal{W}_{3r}$ for all $s > 0$ will eventually hit the boundary of $\mathcal{W}_{3r}$. Let $s_{\mathrm{out}} \in (0, \infty]$ be the largest real number such that

$$\Phi_s(\gamma_i) \in \mathcal{W}_{3r} \setminus \mathcal{W}_r, \qquad \forall s \in (0, s_{\mathrm{out}}).$$

We must have $s_{\mathrm{out}} < b/\delta^2 < \infty$, for

$$0 < \mathcal{S}_{c(L)}(\Phi_{s_{\mathrm{out}}}(\gamma_i)) = \mathcal{S}_{c(L)}(\gamma_i) - \int_0^{s_{\mathrm{out}}} \|\nabla \mathcal{S}_{c(L)}(\Phi_s(\gamma_i))\|^2 < b - \delta^2 s_{\mathrm{out}}.$$

However, since on the time interval $[0, s_{\text{out}}]$ the curve $s \mapsto \Phi_s(\gamma_i)$ crosses a region of width r, we have the estimate

$$\begin{aligned}\mathcal{S}_{c(L)}(\Phi_{s_{\text{out}}}(\gamma_i)) &= \mathcal{S}_{c(L)}(\gamma_i) - \int_0^{s_{\text{out}}} \|\nabla \mathcal{S}_{c(L)}(\Phi_s(\gamma_i))\|^2 \mathrm{d}s \\ &< b - \delta \int_0^{s_{\text{out}}} \|\partial_s \Phi_s(\gamma_i)\| \, \mathrm{d}s \\ &< b - \delta r \\ &< 0,\end{aligned}$$

which contradicts the fact that $\inf \mathcal{S}_{c(L)} = 0$. This completes the proof of the lemma for $e = c(L)$. The general statement follows from this, by observing that

$$\mathcal{S}_e(\Gamma, p) = \mathcal{S}_{c(L)}(\Gamma, p) + (e - c(L))p. \qquad \blacksquare$$

Proof of Theorem 1.3. Let ϵ and $\mathcal{W}$ be the constant and the neighborhood of $\mathcal{K}_1$ given by Lemma 4.2. We require $\mathcal{W}$ to be small enough so that, according to Lemma 2.1, every periodic orbit of the Lagrangian system of L contained in $\mathcal{W}$ is simple. Fix an arbitrary energy value $e \in (c(L), c(L) + \epsilon)$, and consider the anti-gradient flow Φ_s of the free-period action functional $\mathcal{S}_e$ associated to the standard Riemannian metric of $\Lambda S^2 \times (0, \infty)$. We set $s := \inf \mathcal{S}_e|_{\mathcal{W}}$.

We claim that, for each $i \in \mathbb{N}$ large enough, there exists $\gamma_i = (\Gamma_i, p_i) \in \mathcal{W} \cap \{\mathcal{S}_e < s + i^{-1}\}$ such that $\|\nabla \mathcal{S}_e(\gamma_i)\| < i^{-1/2}$. Indeed, assume that this does not hold, which means that there exists $i \in \mathbb{N}$ large enough such that $s + i^{-1} < \inf \mathcal{S}_e|_{\partial \mathcal{W}}$ and $\|\nabla \mathcal{S}_e\| \geq i^{-1/2}$ on $\mathcal{W} \cap \{\mathcal{S}_e < s + i^{-1}\}$. Choose any $\gamma \in \mathcal{W} \cap \{\mathcal{S}_e < s + i^{-1}\}$. Lemma 4.2 readily implies that $\Phi_s(\gamma) \in \mathcal{W}$ for all $s \geq 0$. However, by flowing γ for time $s = 1$ we obtain the contradiction

$$\mathcal{S}_e(\Phi_1(\gamma)) = \mathcal{S}_e(\gamma) - \int_0^1 \|\nabla \mathcal{S}_e(\Phi_s(\gamma))\|^2 \mathrm{d}s < s + i^{-1} - i^{-1} = s.$$

Since $p_i \in [p_{\min}/2, 2p_{\max}]$, and since $\mathcal{S}_e$ satisfies the Palais-Smale condition on $\Lambda S^2 \times [p_{\min}/2, 2p_{\max}]$, γ_i converges to a global minimizer γ of $\mathcal{S}_e|_{\mathcal{W}}$. $\blacksquare$

Proof of Theorem 1.6. Consider an arbitrary energy level $e \in (c(L), c_\infty(L))$. Up to modifying the Lagrangian L away from the sublevel set $E^{-1}(-\infty, e + 1]$, we can assume that L is quadratic at infinity, and in particular its free-period action functional $\mathcal{S}_e$ is well behaved (see Section 2.3).

Since we are looking for infinitely many periodic orbits with energy e, we can assume that the set of critical points $\mathrm{crit}(\mathcal{S}_e)$ is a collection of isolated critical circles. Notice that $\mathcal{S}_e(\gamma) > 0$, since $e > c(L)$. For each $m \in \mathbb{N}$, we introduce the space of paths

$$\mathcal{P}_m := \big\{ P : [0,1] \xrightarrow{C^0} \Lambda S^2 \times (0, \infty) \;\big|\; P(0) \in \gamma,\ P(1) \in \gamma^m \big\},$$

and the minimax value

$$s_m := \inf_{P \in \mathcal{P}_m} \max_{s \in [0,1]} \mathcal{S}_e(P(s)).$$

Since we are working in the 2-sphere, every iterate of γ is still a local minimizer of $\mathcal{S}_e$, see [AMP15, Lemma 4.1]. Therefore

$$s_m > \mathcal{S}_e(\gamma^m) > 0.$$

As $e > c(L)$, the free-period action functional $\mathcal{S}_e$ satisfies the Palais-Smale condition at every non-zero level on the whole $\Lambda S^2 \times (0,\infty)$, see [Abb13, Lemmas 5.1–5.4]. Therefore, s_m is a critical value of the free-period action functional $\mathcal{S}_e$. Actually, a standard deformation argument from critical point theory allows us to find, for each arbitrarily small neighborhood $\mathcal{W} \subset \Lambda S^2 \times (0,\infty)$ of the set of critical points $\mathrm{crit}(\mathcal{S}_e) \cap \mathcal{S}_e^{-1}(s_m)$, a path $P \in \mathcal{P}_m$ such that

$$P([0,1]) \subset \{\mathcal{S}_e < s_m\} \cup \mathcal{W}. \tag{4.1}$$

Assume now by contradiction that the Lagrangian system of L admits only finitely many non-iterated periodic orbits $\gamma_1, ..., \gamma_k$ with energy e. In particular, every such periodic orbit γ_i lies on an isolated critical circle $\mathcal{C}_i$ of $\mathcal{S}_e$. Let us recall the non-mountain pass Theorem for high iterates, which was originally proved in [AMMP17, Theorem 2.6] for magnetic Lagrangians, and extended in [AM16, Lemma 4.3 and proof of Theorem 1.2] to the case of general Tonelli Lagrangians: there exist constants $\overline{m}_i \in \mathbb{N}$ and, for all $m \geq \overline{m}_i$, arbitrarily small open neighborhoods $\mathcal{W}_{i,m}$ of $\psi^m(\mathcal{C}_i)$ such that the inclusion induces an injective map between path-connected components

$$\pi_0(\{\mathcal{S}_e < \mathcal{S}_e(\gamma_i^m)\}) \hookrightarrow \pi_0(\{\mathcal{S}_e < \mathcal{S}_e(\gamma_i^m)\} \cup \mathcal{W}_{i,m}). \tag{4.2}$$

Consider a large enough $m \in \mathbb{N}$ such that $s_m \geq \max\left\{\mathcal{S}_e(\gamma_i^m) \;\middle|\; i = 1, ..., k\right\}$. The set of critical points $\mathrm{crit}(\mathcal{S}_e) \cap \mathcal{S}_e^{-1}(s_e)$ is comprised of finitely many critical circles

$$\mathrm{crit}(\mathcal{S}_e) \cap \mathcal{S}_e^{-1}(s_e) = \psi^{m_1}(\mathcal{C}_{i_1}) \cup ... \cup \psi^{m_k}(\mathcal{C}_{i_k}),$$

where $m_j \geq \overline{m}_{i_j}$ for all $j = 1, ..., k$. We choose the neighborhoods $\mathcal{W}_{i_j,m_j}$ of $\mathcal{C}_{i_j}$ small enough so that they are pairwise disjoint. We set

$$\mathcal{W} := \mathcal{W}_{i_1,m_1} \cup ... \cup \mathcal{W}_{i_k,m_k},$$

and we choose a path $P \in \mathcal{P}_m$ satisfying (4.1). By (4.2), we can modify P in order to obtain a new path $Q \in \mathcal{P}_m$ such that $Q([0,1]) \subset \{\mathcal{S}_e < s_e\}$. This contradicts the definition of the minimax value s_m. ∎

5. A sufficient condition for the inequality $e_0(L) < c_{\mathrm{u}}(L)$

Let M be a closed manifold, and $L : \mathrm{T}M \to \mathbb{R}$ a Tonelli Lagrangian with energy function $E : \mathrm{T}M \to \mathbb{R}$, $E(q,v) = \partial_v L(q,v)v - L(q,v)$. As before, we define

$$e_0(L) = \max_{q \in M} E(q,0)$$

and, for any $e \in \mathbb{R}$, the free-period action functional

$$\mathcal{S}_e : \Lambda M \times (0,\infty) \to \mathbb{R}, \qquad \mathcal{S}_e(\gamma) = \int_0^p \Big[L(\gamma(t), \dot\gamma(t)) + e\Big] \mathrm{d}t.$$

We consider the Mañé critical value of the lift of L to the tangent bundle of the universal cover of M, which is usually denoted by $c_{\mathrm{u}}(L)$. Analogously, $c_{\mathrm{u}}(L)$ is the minimal energy value $e \in \mathbb{R}$ such that $\mathcal{S}_e(\gamma)$ is non-negative for all contractible periodic curves $\gamma = (\Gamma, p) \in \Lambda M \times (0,\infty)$. Since $\mathcal{S}_{e_0(L)}$ vanishes at the constant curves contained in $\{q \in M \mid E(q,0) = e_0(L)\}$, we have that $e_0(L) \leq c_{\mathrm{u}}(L)$. We now give a criterion for the strict inequality $e_0(L) < c_{\mathrm{u}}(L)$.

For all $q \in M$, let us define

$$V(q) = -L(q,0), \qquad \theta_q(v) = \partial_v L(q,0)v, \qquad g_q(v,v) = \partial_{vv}^2 L(q,0)[v,v],$$

where V is a function, θ a one-form, and g a symmetric bilinear form on M. Since L is a Tonelli Lagrangian, g is actually a Riemannian metric on M. Notice that

$$e_0(L) = \max\big(-L(\cdot,0)\big) = \max V.$$

We define the function

$$\lambda : V^{-1}(e_0(L)) \to \mathbb{R}, \qquad \lambda(q) = 2|\mathrm{d}^2V(q)|^{1/2} - |\mathrm{d}\theta_q|, \tag{5.1}$$

where $|\cdot|$ is the norm induced by g, and $\mathrm{d}^2V(q)$ is the Hessian of V at q. The Hessian is well defined, for $V^{-1}(e_0(L)) \subset \mathrm{crit}(V)$.

Proposition 5.1. *If the function λ is somewhere negative, then $e_0(L) < c_{\mathrm{u}}(L)$.*

Proof. Assume that there exists a point $q \in V^{-1}(e_0(L))$ such that $\lambda(q) < 0$, and fix two normal tangent vectors $u, v \in \mathrm{T}_qM$ such that $|u| = |v| = 1$ and

$$\mathrm{d}\theta_q(u,v) = |\mathrm{d}\theta_q|. \tag{5.2}$$

We denote by $D \subset \mathbb{R}^2$ the Euclidean unit disc centered at the origin $\mathbf{0}$, and consider an embedding $\iota : D \hookrightarrow M$ such that $\iota(\mathbf{0}) = q$ and $\mathrm{d}\iota(\mathbf{0})(T_{\mathbf{0}}D) = \mathrm{span}\{u, v\}$. We pull-back L to a Tonelli Lagrangian $L' := L \circ \mathrm{d}\iota : \mathrm{T}D \to \mathbb{R}$ with associated free-period action functionals $\mathcal{S}'_e$, and we set

$$g' := \iota^* g, \qquad \theta' := \iota^*\theta, \qquad V' := V \circ \iota.$$

The analog of the function λ for the pulled-back Lagrangian L' is

$$\lambda' : (V')^{-1}(e_0(L)) \to \mathbb{R}, \qquad \lambda'(x) := 2|\mathrm{d}^2V'(x)|^{1/2} - |\mathrm{d}\theta'_x|,$$

where now $|\cdot|$ denotes the norm associated to the pulled-back metric ι^*g. Notice that $|\mathrm{d}^2V'(\mathbf{0})| \le |\mathrm{d}^2V(q)|$ and, by (5.2), $|\mathrm{d}\theta'_{\mathbf{0}}| = |\mathrm{d}\theta_q|$. Therefore,

$$\lambda'(\mathbf{0}) \le \lambda(q) < 0.$$

Let μ be the Riemannian volume form associated to g' with respect to some orientation on D, and $f : D \to \mathbb{R}$ the function defined by $\mathrm{d}\theta' = f\mu$, so that $|\mathrm{d}\theta'| = |f|$. We fix the orientation on D so that $f(\mathbf{0}) \le 0$. We set $b := |\mathrm{d}^2V'(\mathbf{0})|$, and we fix $a > 0$ small enough such that

$$2\sqrt{a+b} + f(\mathbf{0}) < 0. \tag{5.3}$$

For all $r > 0$ smaller than the injectivity radius of g' at $\mathbf{0}$, we set

$$e_r := e_0(L) + \tfrac{a}{2}r^2,$$

and we denote by γ_r the boundary of the Riemannian ball $B_r(\mathbf{0})$ of g' centered at $\mathbf{0}$ of radius r. We parametrize γ_r counterclockwise with constant speed $|\dot\gamma_r(t)| \equiv s_r$ and period τ_r given by

$$\tau_r := \frac{\ell_r}{r\sqrt{a+b}}, \qquad s_r := r\sqrt{a+b},$$

where ℓ_r is the length of γ_r. Since $\ell_r = 2\pi r + o(r)$, we have

$$\tau_r = \frac{2\pi}{\sqrt{a+b}} + o(1).$$

Using that $e_0(L) = \max V' = V'(\mathbf{0})$, we can estimate

$$L'(x,u) \le -e_0(L) + \theta'_x(u) + \tfrac{1}{2}|u|^2 + \tfrac{1}{2}b\,\mathrm{dist}(\mathbf{0},x)^2 + o(|x|^2) + o(|u|^2).$$

Therefore

$$\begin{aligned} L'(\gamma_r,\dot\gamma_r)+e_r &= \frac{1}{2}s_r^2+\theta'_{\gamma_r}(\dot\gamma_r)+e_r-e_0+\frac{b}{2}r^2+o(r^2)\\ &= \frac{a+b}{2}r^2+\theta'_{\gamma_r}(\dot\gamma_r)+\frac{a+b}{2}r^2+o(r^2)\\ &= (a+b)r^2+\theta'_{\gamma_r}(\dot\gamma_r)+o(r^2). \end{aligned}$$

Moreover

$$\int_0^{\tau_r}\theta'_{\gamma_r(t)}(\dot\gamma_r(t))\mathrm{d}t=\int_{B_r(\mathbf{0})}\mathrm{d}\theta'=\int_{B_r(\mathbf{0})}f\mu=f(\mathbf{0})\pi r^2+o(r^2).$$

Putting together the last two equations, we obtain the estimate

$$\begin{aligned} \mathcal{S}'_{e_r}(\gamma_r) &= \tau_r(a+b)r^2+o(r^2)+f(\mathbf{0})\pi r^2+o(r^2)\\ &= \big(2\sqrt{a+b}+f(\mathbf{0})\big)\pi r^2+o(r^2). \end{aligned}$$

This, together with (5.3), shows that $\mathcal{S}'_{e_r}(\gamma_r)$ is negative for $r>0$ small enough. Therefore, we constructed a periodic curve $\iota\circ\gamma_r$ that is contractible and satisfies $\mathcal{S}_{e_r}(\iota\circ\gamma_r)=\mathcal{S}'_{e_r}(\gamma_r)<0$, which implies that $e_0(L)<e_r<c_{\mathrm{u}}(L)$. ∎

6. Criticality of the Katok metric

In his 1973 seminal paper [Kat73], Katok constructed a non-reversible Finsler metric on every n-sphere with only finitely many closed geodesics. Another description of this Finsler metric, which shows that it is of Randers type, was provided a decade later by Ziller [Zil83]. For the 2-sphere, Ziller's construction goes as follows. Let $g_0=\langle\cdot,\cdot\rangle$ be the round metric on S^2, that is, the metric induced by the embedding of S^2 as the unit sphere of the Euclidean space $\mathbb{R}^3$ with its Euclidean Riemannian metric. We denote by $\psi^t:S^2\to S^2$ the counterclockwise rigid rotation of angle t around the axis passing through the north and south poles of S^2, i.e.

$$\psi^t(x,y,z)=(\cos(t)x-\sin(t)y,\sin(t)x+\cos(t)y,z).$$

Let $W=-y\partial_x+x\partial_y$ be the Killing vector field on (S^2,g) generating this rotation, that is, satisfying $\frac{\mathrm{d}}{\mathrm{d}t}\psi^t=W\circ\psi^t$. We will write $|\cdot|$ for the Riemannian norm induced by g_0 on tangent and cotangent vectors on S^2. We have

$$|W(x,y,z)|=\sqrt{1-z^2}\le 1,\qquad \forall(x,y,z)\in S^2.$$

For $\alpha\in(0,1)$, we consider the Randers metric on cotangent vectors

$$F^*:\mathrm{T}^*S^2\to[0,\infty),\qquad F^*(q,p)=|p|+\alpha\,p(W(q)).$$

The **Katok metric** on S^2 with parameter α is the dual Finsler metric

$$F:\mathrm{T}S^2\to[0,\infty),\qquad F(q,v)=\max_{F^*(q,p)=1}p\,v.$$

Since F^* is of Randers type, the same is true for F. Namely, F has the form

$$F(q,v)=g_q(v,v)^{1/2}+\theta_q(v) \tag{6.1}$$

for some Riemannian metric $g=g_\alpha$ and 1-form $\theta=\theta_\alpha$ on S^2. The Randers metric F admits two distinguished closed geodesics: the ones running clockwise and counterclockwise around the equator. If α is irrational, F has no other closed geodesics. Nevertheless, we have the following theorem.

Theorem 6.1. *The Katok metric $F(q,v) = g_q(v,v)^{1/2} + \theta_q(v)$ with parameter $\alpha \in (0,1)$ is critical (in the sense of Definition 1.9), with*

$$r(g,\theta) = \alpha, \qquad r_\infty(g,\theta) \in \left[\tfrac{2\alpha}{\alpha^2+1}, 1\right].$$

In particular, for each $r \in \left(\alpha, \frac{2\alpha}{\alpha^2+1}\right)$, the Finsler metric

$$F_r(q,v) = r\, g_q(v,v)^{1/2} + \theta_q(v)$$

possesses a waist and infinitely many closed geodesics.

The criticality of Katok's metric is a consequence of the following general result.

Lemma 6.2. *Let $F(q,v) = g_q(v,v)^{1/2} + \theta_q(v)$ be a Randers metric on S^2 such that θ is not exact. Let Z be the vector field on S^2 defined by $g(Z,\cdot) = \theta$, and*

$$N = \left\{ q \in S^2 \;\middle|\; |\theta_q| = \|\theta\|_\infty \right\},$$

where $|\cdot|$ is the norm induced by g, and $\|\cdot\|_\infty$ the associated ∞-norm. If N contains a periodic orbit of the flow of Z, then F is critical.

Proof. Consider the magnetic Lagrangian associated to the Randers metric F

$$L : \mathrm{T}S^2 \to \mathbb{R}, \qquad L(q,v) = \tfrac{1}{2}|v|^2 + \theta_q(v).$$

We already know that $r(g,\theta) = \sqrt{2c(L)} \le \|\theta\|_\infty$. Therefore, all we have to show is that $c(L) \ge \frac{1}{2}\|\theta\|_\infty^2$. Notice that

$$L(q,v) + \tfrac{1}{2}\|\theta\|_\infty^2 = \tfrac{1}{2}\big(|v + Z(q)|^2 + \|\theta\|_\infty^2 - |\theta_q|^2\big).$$

In particular

$$L(q,v) + \tfrac{1}{2}\|\theta\|_\infty^2 = \tfrac{1}{2}|v + Z(q)|^2, \qquad \forall q \in N,\ v \in \mathrm{T}_qS^2. \tag{6.2}$$

Let $\zeta : \mathbb{R}/p\mathbb{Z} \to N$ be a periodic orbit of the flow of Z, and consider the flipped curve $\gamma : \mathbb{R}/p\mathbb{Z} \to N$, $\gamma(t) = \zeta(-t)$, which satisfies $\dot\gamma(t) = -Z(\gamma(t))$. By (6.2), we have

$$\mathcal{S}_{\|\theta\|_\infty^2/2}(\gamma) = \int_0^p \tfrac{1}{2}|\dot\gamma(t) + Z(\gamma(t))|^2 \mathrm{d}t = 0,$$

which implies the desired inequality $\|\theta\|_\infty^2/2 \ge c(L)$. ∎

Remark 6.3. The hypotheses of Lemma 6.2 are satisfied, for instance, if the pair (g,θ) is rotationally symmetric, namely if there exist spherical coordinates $(\phi,\tau) \in [0,\pi] \times \mathbb{R}/2\pi\mathbb{Z}$ on S^2 such that

$$g = a(\phi)^2\mathrm{d}\phi^2 + b(\phi)^2\mathrm{d}\tau^2, \qquad \theta = c(\phi)\mathrm{d}\tau$$

for some functions $a, b, c : [0,\pi] \to \mathbb{R}$ with c not identically zero. ∎

Proof of Theorem 6.1. If $F^* : \mathrm{T}^*M \to [0,\infty)$, $F^*(q,p) = |p| + \alpha\, p(W(q))$ is a cotangent Randers metric on an arbitrary manifold M, then the the dual metric $F : \mathrm{T}M \to [0,\infty)$ is defined by the equation

$$\big|F(q,v)^{-1}v - \alpha W(q)\big|^2 = 1.$$

Therefore, F has the form $F(q,v) = g(v,v)^{1/2} + \theta_q(v)$, where the Riemannian metric g and the 1-form θ on M are given by

$$g_q(v,v)^{1/2} = \frac{\sqrt{|v|^2(1-\alpha^2|W(q)|^2) - \alpha^2\langle v, W(q)\rangle^2}}{1-\alpha^2|W(q)|^2},$$
$$\theta_q(v) = -\frac{\alpha\langle v, W(q)\rangle}{1-\alpha^2|W(q)|^2},$$

see also [Ben16, Eq. (3.13)]. Now, if $M = S^2$ and F is the Katok metric with parameter α, for each $q = (x,y,z) \in S^2$ and $v = v_x\partial_x + v_y\partial_y + v_z\partial_z \in \mathrm{T}_qS^2$, we have

$$g_q(v,v)^{1/2} = \frac{\sqrt{|v|^2(1-\alpha^2(1-z^2)) - \alpha^2(xv_y - yv_x)^2}}{1-\alpha^2(1-z^2)},$$
$$\theta_q(v) = -\frac{\alpha\,(xv_y - yv_x)}{1-\alpha^2(1-z^2)}$$

It is convenient to rewrite these expressions in spherical coordinates on S^2. We denote by $\phi \in [0,\pi]$ and $\tau \in \mathbb{R}/2\pi\mathbb{Z}$ the polar angle and the azimuthal angle respectively, as in Figure 1. For each $q = (\phi,\tau) \in [0,\pi]\times\mathbb{R}/2\pi\mathbb{Z}$ and $v = (v_\phi, v_\tau) \in \mathbb{R}^2$, we have

$$g_q(v,v) = \frac{1}{1-\alpha^2\sin^2\phi}v_\phi^2 + \Big(\frac{\sin\phi}{1-\alpha^2\sin^2\phi}\Big)^2 v_\tau^2,$$
$$\theta_q(v) = -\frac{\alpha\sin^2\phi}{1-\alpha^2\sin^2\phi}v_\tau.$$

In order to simplify the notation, we will denote by $|\cdot|$ the norm on tangent and cotangent vectors induced by g, and by $\|\cdot\|_\infty$ the associated L^∞ norm on 1-forms. With this notation, we have

$$|\theta_{(\phi,\tau)}| = \alpha\,\sin\phi,$$
$$\|\theta\|_\infty = \alpha.$$

Therefore

$$N := \big\{(\phi,\tau) \in [0,\pi]\times\mathbb{R}/2\pi\mathbb{Z} \;\big|\; |\theta_q| = \|\theta\|_\infty\big\} = \{\pi/2\}\times\mathbb{R}/2\pi\mathbb{Z},$$

that is, the subset N from Lemma 6.2 is the equator oS^2. The dual vector field of θ with respect to g is given by $Z = -\alpha(1-\alpha^2\sin^2\phi)\partial_\tau$. In particular, N is a periodic orbit of Z. Lemma 6.2 implies that the Katok metric F is critical, and therefore

$$r(g,\theta) = \|\theta\|_\infty = \alpha.$$

The curve that winds counterclockwise around the equator of S^2 is given in spherical coordinates by

$$\gamma_0 : \mathbb{R}/\mathbb{Z} \to N \subset [0,\pi]\times\mathbb{R}/2\pi\mathbb{Z}, \qquad \gamma_0(t) = (\pi/2, 2\pi t).$$



FIGURE 1. Spherical coordinates on S^2.

We claim that γ_0 is a waist of the Finsler metric F_r, for each $r \in \left(\alpha, \frac{2\alpha}{\alpha^2+1}\right)$. Indeed, the length functional associated to F_r can be written in spherical coordinates as

$$\mathcal{L}_r : C^\infty(\mathbb{R}/\mathbb{Z}, [0,\pi] \times \mathbb{R}/2\pi\mathbb{Z}) \to \mathbb{R},$$

$$\mathcal{L}_r(\gamma) = \int_0^1 \left[r\left(\frac{(\dot\phi)^2}{1-\alpha^2\sin^2\phi} + \left(\frac{\dot\tau\,\sin\phi}{1-\alpha^2\sin^2\phi} \right)^2 \right)^{1/2} - \frac{\alpha\,\sin^2\phi}{1-\alpha^2\sin^2\phi}\dot\tau \right] \mathrm{d}t,$$

where we have written $\gamma = (\phi, \tau)$. If γ is sufficiently C^1-close to γ_0, then $\dot\tau(t) > 0$ for all $t \in \mathbb{R}/\mathbb{Z}$, and we can estimate

$$\mathcal{L}_r(\gamma) \geq \int_0^1 \underbrace{\frac{r\,\sin\phi(t) - \alpha\,\sin^2\phi(t)}{1-\alpha^2\sin^2\phi(t)}}_{=: f_{r,\alpha}(\phi(t))} \dot\tau(t)\,\mathrm{d}t.$$

The function $f_{r,\alpha} : [0,\pi] \to \mathbb{R}$ has a local minimum at $\pi/2$ if and only if $r < \frac{2\alpha}{\alpha^2+1}$. Therefore, for each $r \in \left(\alpha, \frac{2\alpha}{\alpha^2+1}\right)$, we can further estimate

$$\mathcal{L}_r(\gamma) \geq f_{r,\alpha}(\pi/2) \int_0^1 \dot\tau(t)\mathrm{d}t = \frac{r-\alpha}{1-\alpha^2} 2\pi = \mathcal{L}_r(\gamma_0).$$

This shows that γ_0 is a waist of $\mathcal{L}_r$, which implies that $r_\infty(g,\theta) \geq \frac{2\alpha}{\alpha^2+1}$. This, together with Theorem 1.8, implies the existence of infinitely many closed geodesics with respect to the Randers metric F_r, for each $r \in \left(\alpha, \frac{2\alpha}{\alpha^2+1}\right)$. ∎

REFERENCES

[AAB+17] A. Abbondandolo, L. Asselle, G. Benedetti, M. Mazzucchelli, and I. A. Taimanov, *The multiplicity problem for periodic orbits of magnetic flows on the 2-sphere*, Adv. Nonlinear Stud. **17** (2017), no. 1, 17–30.

[Abb13] A. Abbondandolo, *Lectures on the free period Lagrangian action functional*, J. Fixed Point Theory Appl. **13** (2013), no. 2, 397–430.

[AM16] L. Asselle and M. Mazzucchelli, *On Tonelli periodic orbits with low energy on surfaces*, arXiv:1601.06692, to appear in Trans. Amer. Math. Soc., 2016.

[AMMP17] A. Abbondandolo, L. Macarini, M. Mazzucchelli, and G. P. Paternain, *Infinitely many periodic orbits of exact magnetic flows on surfaces for almost every subcritical energy level*, J. Eur. Math. Soc. **19** (2017), 551–579.

[AMP15] A. Abbondandolo, L. Macarini, and G. P. Paternain, *On the existence of three closed magnetic geodesics for subcritical energies*, Comment. Math. Helv. **90** (2015), no. 1, 155–193.

[Ben16] G. Benedetti, *Magnetic Katok examples on the two-sphere*, Bull. Lond. Math. Soc. **48** (2016), no. 5, 855–865.

[CI99] G. Contreras and R. Iturriaga, *Global minimizers of autonomous Lagrangians*, 22º Colóquio Brasileiro de Matemática, IMPA, Rio de Janeiro, 1999.

[CIPP98] G. Contreras, R. Iturriaga, G. P. Paternain, and M. Paternain, *Lagrangian graphs, minimizing measures and Mañé's critical values*, Geom. Funct. Anal. **8** (1998), no. 5, 788–809.

[CIPP00] ———, *The Palais-Smale condition and Mañé's critical values*, Ann. Henri Poincaré **1** (2000), no. 4, 655–684.

[CMP04] G. Contreras, L. Macarini, and G. P. Paternain, *Periodic orbits for exact magnetic flows on surfaces*, Int. Math. Res. Not. (2004), no. 8, 361–387.

[Fat08] A. Fathi, *Weak KAM theorem in Lagrangian dynamics*, Cambridge Univ. Press, forthcoming, preliminary version number 10, 2008.

[Kat73] A. B. Katok, *Ergodic perturbations of degenerate integrable Hamiltonian systems*, Izv. Akad. Nauk SSSR Ser. Mat. **37** (1973), 539–576.

[Mañ97] R. Mañé, *Lagrangian flows: the dynamics of globally minimizing orbits*, Bol. Soc. Brasil. Mat. (N.S.) **28** (1997), no. 2, 141–153.

[Mat91] J. N. Mather, *Action minimizing invariant measures for positive definite Lagrangian systems*, Math. Z. **207** (1991), no. 2, 169–207.

[Pat99] G. P. Paternain, *On two noteworthy deformations of negatively curved riemannian metrics*, Discrete Contin. Dynam. Systems **5** (1999), no. 3, 639–650.

[Sor15] A. Sorrentino, *Action-minimizing methods in Hamiltonian dynamics: an introduction to Aubry-Mather theory*, Mathematical Notes, vol. 50, Princeton University Press, Princeton, NJ, 2015.

[Tai91] I. A. Taimanov, *Non-self-itersecting closed extremals of multivalued or not everywhere positive functionals*, Izv. Akad. Nauk SSSR Ser. Mat. **55** (1991), no. 2, 367–383.

[Tai92] ———, *Closed non-self-intersecting extremals of multivalued functionals*, Sibirsk. Mat. Zh. **33** (1992), no. 4, 155–162, 223.

[Zil83] W. Ziller, *Geometry of the Katok examples*, Ergodic Theory Dynam. Systems **3** (1983), no. 1, 135–157.

Gabriele Benedetti
Universität Leipzig, Mathematisches Institut
Augustusplatz 10, D-04109 Leipzig, Germany
E-mail address: gabriele.benedetti@math.uni-leipzig.de

Marco Mazzucchelli
CNRS, École Normale Supérieure de Lyon, UMPA
46 allée d'Italie, 69364 Lyon Cedex 07, France
E-mail address: marco.mazzucchelli@ens-lyon.fr